

Nonlinear gauge transformation for a class of Schrödinger equations containing complex nonlinearities

G. Kaniadakis* and A.M. Scarfone†

*Dipartimento di Fisica - Politecnico di Torino - Corso Duca degli Abruzzi 24, 10129 Torino, Italy;
Istituto Nazionale di Fisica della Materia - Unità del Politecnico di Torino,
Corso Duca degli Abruzzi 24, 10129 Torino, Italy*

(Dated: February 1, 2008)

We consider a wide class of nonlinear canonical quantum systems described by a one-particle Schrödinger equation containing a complex nonlinearity. We introduce a nonlinear unitary transformation which permits us to linearize the continuity equation. In this way we are able to obtain a new quantum system obeying to a nonlinear Schrödinger equation with a real nonlinearity. As an application of this theory we consider a few already studied Schrödinger equations as that containing the nonlinearity introduced by the exclusion-inclusion principle, the Doebner-Goldin equation and others.

PACS numbers: PACS numbers: 03.65.-w, 11.15.-q

Let us introduce the density of Lagrangian

$$\mathcal{L} = i \frac{\hbar}{2} \left(\psi^* \frac{\partial \psi}{\partial t} - \psi \frac{\partial \psi^*}{\partial t} \right) - \frac{\hbar^2}{2m} \left| \frac{\partial \psi}{\partial x} \right|^2 - U[\psi, \psi^*], \quad (1)$$

which describes a class of one dimensional nonrelativistic and canonical quantum systems. In Eq. (1) the nonlinear real potential $U[\psi, \psi^*]$ is a functional of the fields ψ and ψ^* . We make use of the notation $U[a]$ to indicate that U is a functional of the field a and of its spatial derivative, that is:

$$U[a] = U(a_0, a_1, a_2, a_3, \dots), \quad (2)$$

with $a_n = \partial^n a / \partial x^n$ and $a_0 \equiv a$.

We will use also the hydrodynamic fields $\rho(x, t)$ and $S(x, t)$ defined as:

$$\rho = \psi \psi^*, \quad (3)$$

$$S = i \frac{\hbar}{2} \log \left(\frac{\psi^*}{\psi} \right), \quad (4)$$

which represent respectively the modulo and the phase of the field ψ :

$$\psi(x, t) = \rho^{1/2}(x, t) \exp \left[\frac{i}{\hbar} S(x, t) \right]. \quad (5)$$

The evolution equations of the fields $a \equiv \psi, \psi^*, \rho, S$ can be obtained from the action of the system:

$$\mathcal{A} = \int \mathcal{L} dx dt, \quad (6)$$

by using the minimal action principle:

$$\frac{\delta \mathcal{A}}{\delta a} = 0. \quad (7)$$

Taking into account that the functional derivative of the potential term in the action $U[a]$ is defined as [1]:

$$\frac{\delta}{\delta a} \int U[a] dx dt = \sum_{n=0} (-1)^n \frac{\partial^n}{\partial x^n} \left(\frac{\partial}{\partial a_n} U[a] \right), \quad (8)$$

it is easy to see that the evolution equation of the field ψ can be obtained from (7), posing $a \equiv \psi^*$ and is given by the following Schrödinger equation which contains a complex nonlinearity:

$$i \hbar \frac{\partial \psi}{\partial t} = - \frac{\hbar^2}{2m} \frac{\partial^2 \psi}{\partial x^2} + W[\rho, S] \psi + i \mathcal{W}[\rho, S] \psi. \quad (9)$$

The real $W[\rho, S]$ and the imaginary $\mathcal{W}[\rho, S]$ part are given by the following expressions:

$$W[\rho, S] = \frac{\delta}{\delta \rho} \int U[\rho, S] dx dt, \quad (10)$$

$$\mathcal{W}[\rho, S] = \frac{\hbar}{2\rho} \frac{\delta}{\delta S} \int U[\rho, S] dx dt. \quad (11)$$

The evolution equation for the field ρ is obtained directly from (7), posing $a \equiv S$ (now S is the field canonically conjugated to the field $-\rho$). We obtain the equation:

$$\frac{\partial \rho}{\partial t} + \frac{\partial j_\psi}{\partial x} = \frac{\partial}{\partial S} U[\rho, S], \quad (12)$$

where the quantum current j_ψ takes the expression:

$$j_\psi = \frac{S_1}{m} \rho + \sum_{n=0} (-1)^n \frac{\partial^n}{\partial x^n} \left(\frac{\partial}{\partial S_{n+1}} U[\rho, S] \right). \quad (13)$$

We remark that Eq. (12) is the continuity equation and the term in the right hand side represents a source for the field ρ . When the conservation of the number of particles $N = \int \rho dx$ is required, the hypothesis that the potential $U[\rho, S]$ does not depend on S but only on its derivative must be introduced, therefore the Eq. (12) takes the form:

$$\frac{\partial \rho}{\partial t} + \frac{\partial j_\psi}{\partial x} = 0, \quad (14)$$

We note that Eq. (14) can be obtained directly from (9) and from its complex conjugate, performing the standard

procedure. We can see that the imaginary part $\mathcal{W}(\rho, S)$ is responsible for the nonlinearity of the expression of the current j_ψ (13).

Let us introduce the following transformation for the field ψ :

$$\psi(x, t) \rightarrow \phi(x, t) = \mathcal{U}[\psi, \psi^*] \psi(x, t), \quad (15)$$

which allows to eliminate the imaginary part of the evolution equation of the field ψ , which corresponds also to linearize the expression of the current j_ψ .

The operator $\mathcal{U}$, generating this transformation, is unitary $\mathcal{U}^\dagger = \mathcal{U}^{-1}$ and is defined by:

$$\begin{aligned} \mathcal{U}[\psi, \psi^*] = & \exp \left[i \frac{m}{\hbar} \sum_{n=0} (-1)^n \right. \\ & \times \left. \int \frac{1}{\rho} \frac{\partial^n}{\partial x^n} \left(\frac{\partial}{\partial S_{n+1}} U[\rho, S] \right) dx \right]. \end{aligned} \quad (16)$$

If we write the field ϕ in terms of the hydrodynamic fields ρ, σ :

$$\phi(x, t) = \rho^{1/2}(x, t) \exp \left[\frac{i}{\hbar} \sigma(x, t) \right], \quad (17)$$

and, due to the unitarity of the transformation, the modulo of ϕ is equal to the modulo of the field ψ , while the phase σ is given by:

$$\sigma = S + \frac{m}{\hbar} \sum_{n=0} (-1)^n \int \frac{1}{\rho} \frac{\partial^n}{\partial x^n} \left(\frac{\partial}{\partial S_{n+1}} U[\rho, S] \right) dx. \quad (18)$$

By accepting the statement made by Feynman and Hibbs ([2], p.96): "*Indeed all measurements of quantum-mechanical systems could be made to reduce eventually to position and time measurements*", (see also [3]) the two wave functions ψ and ϕ represent the same physical system and, as a consequence, we can interpret the Eq. (15) as a nonlinear gauge transformation of the function described by Eq. (9).

From Eq. (9) and taking into account the transformation (15), it is easy to obtain the following evolution equation for the field ϕ :

$$\begin{aligned} i \hbar \frac{\partial \phi}{\partial t} = & -\frac{\hbar^2}{2m} \frac{\partial^2 \phi}{\partial x^2} + W[\rho, S] \phi \\ & -\frac{1}{2} m \left[\sum_{n=0} (-1)^n \frac{1}{\rho} \frac{\partial^n}{\partial x^n} \left(\frac{\partial U[\rho, S]}{\partial S_{n+1}} \right) \right]^2 \phi \\ & -m \sum_{n=0} (-1)^n \frac{\partial}{\partial t} \left[\int \frac{1}{\rho} \frac{\partial^n}{\partial x^n} \left(\frac{\partial U[\rho, S]}{\partial S_{n+1}} \right) dx \right] \phi \\ & - \sum_{n=0} (-1)^n \frac{S_1}{\rho} \frac{\partial^n}{\partial x^n} \left(\frac{\partial U[\rho, S]}{\partial S_{n+1}} \right) \phi. \end{aligned} \quad (19)$$

Note that the nonlinearity appearing in Eq. (19) is now real. The continuity equation of the system takes the

form:

$$\frac{\partial \rho}{\partial t} + \frac{\partial j_\phi}{\partial x} = 0 \quad (20)$$

where the current j_ϕ has the standard expression of the linear quantum mechanics:

$$j_\phi = \frac{\sigma_1}{m} \rho. \quad (21)$$

The gauge transformation (15) and (16) makes real the complex nonlinearity in the evolution equation, and makes non canonical the new dynamical system. However, this transformation may be useful to describe the evolution of system by means of an equation containing a real nonlinearity. We note that nonlinear transformations have been introduced and used systematically for the first time in order to study nonlinear Schrödinger equations as the Doebner-Goldin one in Ref. [3].

Let us apply the proposed transformation to a few already known equations describing systems of collectively interacting particles.

The first example is given by the canonical Doebner-Goldin equation ($c_1 = -c_4 = 1, c_2 = c_3 = c_5 = 0$) [3, 4] that can be obtained from (1) where the potential $U[\rho, S]$ has the following form:

$$U[\rho, S] = \frac{D}{2} (\rho_1 S_1 - \rho S_2). \quad (22)$$

A complex nonlinearity is generated in the evolution equation of the field ψ , with real and imaginary part given respectively by:

$$W[\rho, S] = -m D \frac{\partial}{\partial x} \left(\frac{j_\psi}{\rho} \right), \quad (23)$$

$$\mathcal{W}[\rho, S] = \frac{\hbar D}{2\rho} \frac{\partial^2 \rho}{\partial x^2}. \quad (24)$$

The quantum current j_ψ takes the form of a Fokker-Planck current:

$$j_\psi = \frac{S_1}{m} \rho + D \rho_1, \quad (25)$$

resulting to be the sum of two terms, the former is a drift current while the latter is a Fick current. The generator of the transformation $\mathcal{U}$ (16) takes the form:

$$\mathcal{U}[\psi, \psi^*] = \exp \left(\frac{i}{\hbar} m D \log \rho \right). \quad (26)$$

This is a particular case of a class of transformations introduced by Doebner and Goldin and permits to write the evolution equation:

$$i \hbar \frac{\partial \phi}{\partial t} = -\frac{\hbar^2}{2m} \frac{\partial^2 \phi}{\partial x^2} + 2m D^2 \frac{1}{\rho^{1/2}} \frac{\partial^2 \rho^{1/2}}{\partial x^2} \phi. \quad (27)$$

Equation (27) was studied in Ref. [5]; after rescaling the phase: $\sigma \rightarrow \sqrt{1 - (2mD/\hbar)^2} \sigma$, it reduces to the linear

Schrödinger equation.

As a second example we consider a nonlinear Schrödinger equation, introduced in Ref. [6], where the complex nonlinearity is originated by the potential:

$$U[\rho, S] = \frac{\hbar^2 \lambda^2}{8m} \rho^3 - \frac{\hbar \lambda}{2m} \rho^2 S_1, \quad (28)$$

and its real and imaginary part take respectively the form:

$$W[\rho, S] = \frac{3\hbar^2 \lambda^2}{8m} \rho^2 - \frac{\hbar \lambda}{m} \rho S_1 \quad (29)$$

$$\mathcal{W}[\rho, S] = \frac{\hbar^2 \lambda}{2m} \rho_1. \quad (30)$$

Within this model the quantum current is given by:

$$j_\psi = \frac{S_1}{m} \rho - \frac{\hbar \lambda}{2m} \rho^2. \quad (31)$$

The generator $\mathcal{U}$ of the transformation (16) takes the simple form:

$$\mathcal{U}[\psi, \psi^*] = \exp\left(-\frac{i\lambda}{2} \int \rho dx\right), \quad (32)$$

while the evolution equation for the new complex quantity ϕ , containing a real nonlinearity takes the form:

$$i\hbar \frac{\partial \phi}{\partial t} = -\frac{\hbar^2}{2m} \frac{\partial^2 \phi}{\partial x^2} - \lambda \hbar j_\phi \phi. \quad (33)$$

The last application is given by a nonlinear Schrödinger equation, proposed recently by us [7, 8], describing a system of particles obeying to an exclusion-inclusion principle (EIP), which are originated from a collective interaction. The potential $U[\rho, S]$ in this model is given by:

$$U[\rho, S] = \kappa \frac{(S_1 \rho)^2}{2m}. \quad (34)$$

The real and imaginary part of the nonlinearity in the evolution equation are:

$$W[\rho, S] = \kappa \frac{m}{\rho} \left(\frac{j_\psi}{1 + \kappa \rho} \right)^2, \quad (35)$$

$$\mathcal{W}[\rho, S] = -\kappa \frac{\hbar}{2\rho} \frac{\partial}{\partial x} \left(\frac{j_\psi \rho}{1 + \kappa \rho} \right). \quad (36)$$

while the current j_ψ is given by:

$$j_\psi = \frac{S_1}{m} \rho (1 + \kappa \rho). \quad (37)$$

The parameter κ quantifies the EIP, which acts as an exclusion principle (inclusion) if $\kappa < 0$ ($\kappa > 0$), since the factor $1 + \kappa \rho$, appearing in the expression of j_ψ becomes an inhibition (enhancement) factor. The gauge transformation $\mathcal{U}$ is:

$$\mathcal{U}[\psi, \psi^*] = \exp\left(\frac{i\kappa}{\hbar} \int \rho S_1 dx\right), \quad (38)$$

and the evolution equation for the field ϕ takes the form:

$$i\hbar \frac{\partial \phi}{\partial t} = -\frac{\hbar^2}{2m} \frac{\partial^2 \phi}{\partial x^2} + \kappa m \frac{j_\phi^2}{\rho(1 + \kappa \rho)} \phi - \kappa \frac{\hbar^2}{4m} \left[\frac{\partial^2 \rho}{\partial x^2} - \frac{1}{\rho} \left(\frac{\partial \rho}{\partial x} \right)^2 \right] \phi. \quad (39)$$

As a conclusion, it is worth to remark that Eq. (16) defines a wide class of nonlinear gauge transformations and allows to reduce the nonlinearity in the evolution equation from complex to real, in several physical systems.

* Electronic address: kaniadakis@polito.it

† Electronic address: scarfone@polito.it

- [1] P. J. Olver: *Applications of Lie Groups to Differential Equations*, Springer, New York, 1986.
- [2] R. P. Feynman and A. R. Hibbs: *Quantum Mechanics and Path Integrals*, McGraw-Hill, New York, 1965.
- [3] H. -D. Doebner and G. A. Goldin: *Phys. Rev. A* **54** (1996), 3764-3771.
- [4] H. -D. Doebner, G. A. Goldin: *J. Phys. A* **27** (1994) 1771-1780.
- [5] F. Guerra and M. Pusterla: *Lett. Nuovo Cimento* **34** (1982), 351-356.
- [6] R. Jackiw: *A Nonrelativistic Chiral Soliton in One Dimension*, preprint MIT-CTP N2587, hep-th/9611185 (1996).
- [7] G. Kaniadakis, P. Quarati, and A. M. Scarfone: *Physica A* **255** (1998), 474-482.
- [8] G. Kaniadakis, P. Quarati and A. M. Scarfone: *Phys. Rev. E* **58** (1998), 5574-5585.